

ToxiBan®

Advantages

Activated charcoal is generally called the “universal antidote” by toxicologists, since it is active in adsorbing most toxicants except inorganic salts and heavy metals. ToxiBan is the “gold standard” for treating animal poisoning.

- Available in three formulations (granules, suspension and suspension with Sorbitol)
- MedChar™, activated charcoal, USP, adsorbs toxins in the gastrointestinal tract
- Kaolin adds extra adsorbing qualities while protecting the intestinal lining
- Sorbitol is a mild hyperosmotic cathartic

Vet-A-Mix offers a complete line of quality, research-proven nutritional supplements and drug dosage forms.



**Advanced
activated
charcoal for
effective
adsorption
of ingested
poisons**



LLOYD, Inc.
800-831-0004
www.lloydinc.com

ToxiBan

Toxicants That Bind to MedChar™, Activated Charcoal, USP

Acetaminophen	Diazepam	Methyl salicylate	Phenylbutazone
Amphetamines	2,4-D (dichlorophenoxy acetic acid)	Morphine	Phenylpropanolamine hydrochloride
Anthelmintics	Digitalis	Muscarine	Phenytoin
Antibiotics	Digitoxin	Narcotics	Quinacrine hydrochloride
Aspirin	Ethylene glycol*	Nicotine	Quinidine
Atropine	Fungicides	Nortriptyline	Quinine
Barbiturates	Hexachlorophene	Organic iodine	Rodenticides
Camphor	Ipecac	Organic metal compounds	Salicylamide
Cantharides	Isoniazid	Organochlorine insecticides	Salicylates
Carbamazepine	Malathion	Organophosphorus insecticides	Strychnine
Carbamates	Mefenamic acid	Parathion	Sulfonamides
Chlordane	Meprobamate	Phenobarbitol	Theophylline
Chloroquine	Methylene blue	Phenothiazine	Tricyclic antidepressants
Chlorpheniramine			
Cocaine			

*Tests for ethylene glycol must be conducted before ToxiBan is administered to prevent false positive reactions.

Indications and Usage:

ToxiBan® is most effective when administered as soon as ingestion of a toxicant is suspected. It may be used in some emergencies when absorption of the toxicant is nearly complete or the exposure was via a parenteral route. This latter application usually involves repetitive or multiple doses. Multiple doses may be used to interrupt the enterohepatic circulation of some toxins, thereby preventing their reabsorption. See package insert for more detailed information.

Contraindications:

There are no known absolute contraindications to the use of activated charcoal. It should not be given simultaneously or shortly before the oral administration of other therapeutic agents such as antibiotics, vitamins or amino acids. Antibiotic therapy should be administered parenterally when ToxiBan is used.

Dosage:

Suspension and Suspension w/Sorbitol

For small animals, use 10 to 20 mL per kg (5 to 10 mL per pound) body weight. For large animals, use 4 to 12 mL per kg (2 to 6 mL per pound) body weight.

Granules

To make a thin suspension, mix 1 volume measure with 5 to 7 parts cold water (Example: 1 level cup ToxiBan to 6 cups water). Agitate vigorously for 10 to 30 seconds. For small animals, use 2 to 4 grams per kg (1 to 2 grams per pound) body weight. For large animals, use 0.75 to 2.0 grams per kilogram (0.35 to 0.9 grams per pound). One pound (453.6 grams) normally treats an animal weighing 225 to 600 kilograms (500 to 1300 pounds).

Administration:

ToxiBan Suspension and Suspension with Sorbitol may be given directly or mixed with a small amount of cold water to reduce viscosity. For Granule administration, use as a powder in food or dilute to create a suspension. When a suspension is created, agitate frequently to keep liquid uniform. A stomach tube is preferred, but an oral drench may be used in emergency.

Precautions:

ToxiBan is an adjunct in the management of poisoning emergencies. Prior to use, proper life support and gastric emptying must be implemented, if indicated. Because more than an hour may pass before ToxiBan can be administered without regurgitation after an emetic has been given, it may be more effective to give ToxiBan immediately because adsorption of a toxicant may be a more effective removal than vomiting.

ToxiBan produces black stools. These stools may have a diarrhea consistency and may persist for several hours. Sorbitol can create a profound cathartic effect. Proper attention should be provided to fluid and electrolyte needs to prevent dehydration, possible hypernatremia and even hypotension. ToxiBan Suspension should be used for multiple use, and cathartics should only be used intermittently during multiple dose activated charcoal use.

Ingredients:

MedChar™ (activated charcoal, USP), Kaolin, and Sorbitol (in granules and suspension with sorbitol). Suspensions also contain suspending agents and preservatives. Granules contain wetting and dispersing agents.

How Supplied:

Suspensions

240 mL bottles, 12 per carton.....List No. 1151
(w/Sorbitol).....List No. 1155

Granules

454 g (1 pound) bottles, 12 per carton.....List No. 0423
5 kg (11 pound) pail.....List No. 0422

KEEP OUT OF REACH OF CHILDREN



P. O. Box 130 • Shenandoah, Iowa 51601 U.S.A.
800-831-0004 • www.lloydinc.com