



# Butorphic®

Butorphanol tartrate sterile solution  
10 mg/mL

Opiate agonist-antagonist for 3-4 hour pain relief.

Only Butorphic Provides the Convenience of a 20 mL Vial

- Lower Inventory cost
- Less Chance of Vial Puncture Contamination
- Fewer Expiry Worries
- Less Time Exposed to Extreme Field Storage Conditions
- Unparalleled Dosing Economy



Butorphanol tartrate, opiate agonist-antagonist, has long been a staple in veterinary medicine for fast acting relief of moderate to severe pain.

The Butorphic brand provides you the quality clinical performance you demand along with the unprecedented economic advantage you have come to expect from LLOYD, Inc.



1-800-831-0004  
[www.lloydinc.com](http://www.lloydinc.com)

# Butorphic® INJECTION

Butorphanol Tartrate  
Veterinary Injection **IV**


## CAUTION

Federal law restricts this drug to use by or on the order of a licensed veterinarian.

## DESCRIPTION

Butorphic Injection (butorphanol tartrate) is a totally synthetic, centrally acting, narcotic agonist-antagonist analgesic with potent antitussive activity. It is a member of the phenanthrene series. The chemical name is Morphinan-3, 14-diol, 17-(cyclobutylmethyl)-, (-)-, (S-(R\*, R\*))2,3-dihydroxybutanedioate (1:1)(salt). It is a white, crystalline, water soluble substance having a molecular weight of 477.55; its molecular formula is  $C_{21}H_{29}NO_2 \cdot C_4H_6O_6$ .

## Chemical Structure:



Each mL of Butorphic Injection contains 10 mg butorphanol base (as tartrate), 3.3 mg citric acid, USP, 6.4 mg sodium citrate, USP, 4.7 mg sodium chloride, USP, and 0.1 mg benzethonium chloride, USP, q.s. with water for injection, USP.

## CLINICAL PHARMACOLOGY

### Comparative Pharmacology

In animals, butorphanol has been demonstrated to be 4 to 30 times more potent than morphine and pentazocine (Talwin®-V) respectively.<sup>1</sup> In humans, butorphanol has been shown to have 5 to 7 times the analgesic activity of morphine and 20 times that of pentazocine.<sup>2,3</sup> Butorphanol has 15 to 20 times the oral antitussive activity of codeine or dextromethorphan in dogs and guinea pigs.<sup>4</sup>

As an antagonist, butorphanol is approximately equivalent to nalorphine and 30 times more potent than pentazocine.<sup>1</sup>

Cardiopulmonary depressant effects are minimal after treatment with butorphanol as demonstrated in dogs<sup>5</sup>, humans<sup>6,7</sup> and horses.<sup>9</sup> Unlike classical narcotic agonist analgesics which are associated with decreases in blood pressure, reduction in heart rate, and concomitant release of histamine, butorphanol does not cause histamine release.<sup>1</sup> Furthermore, the cardiopulmonary effects of butorphanol are not distinctly dosage related but rather reach a ceiling effect beyond which further dosage increases result in relatively lesser effects.


Reproduction: Studies performed in mice and rabbits revealed no evidence of impaired fertility or harm to the fetus due to butorphanol tartrate. In the female rat, parenteral administration was associated with increased nervousness and decreased care for the newborn, resulting in a decreased survival rate of the newborn. This nervousness was seen only in the rat species.

## Equine Pharmacology

Following intravenous injection in horses, butorphanol is largely eliminated from the blood within 3 to 4 hours. The drug is extensively metabolized in the liver and excreted in the urine.

In ponies butorphanol given intramuscularly at a dosage of 0.22 mg/kg was shown to alleviate experimentally induced visceral pain for about 4 hours.<sup>9</sup>

In horses, intravenous dosages of butorphanol ranging from 0.05 to 0.4 mg/kg were shown to be effective in alleviating visceral and superficial pain for at least four hours, as illustrated in the following figure:



\*Pain threshold in butorphanol-treated colicky horses relative to placebo controls.

A definite dosage-response relationship was detected in that butorphanol dosage of 0.1 mg/kg was more effective than 0.05 mg/kg but not different from 0.2 mg/kg in alleviating deep abdominal pain.

## Acute Equine Studies

Rapid intravenous administration of butorphanol at a dosage of 2.0 mg/kg (20 times the recommended dosage) to a previously unmedicated horse resulted in a brief episode of inability to stand, muscle fasciculation, a convulsive seizure of 6 seconds duration, and recovery within three minutes. The same dosage administered after 10 successive daily 1.0 mg/kg dosages of butorphanol resulted only in transient sedative effects. During the 10 day course of administration at 1.0 mg/kg (10 times the recommended use level) in two horses, the only detectable drug effects were transient behavioral changes typical of narcotic agonist activity. These included muscle fasciculation about the head and neck, dysphoria, lateral nystagmus, ataxia, and salivation. Repeated administration of butorphanol at 1.0 mg/kg (10 times the recommended dose) every four hours for 48 hours caused constipation in one of two horses.

## Subacute Equine Studies

Horses were found to tolerate butorphanol given intravenously at dosages of 0.1, 0.3, and 0.5 mg/kg every 4 hours for 48 hours followed by once daily injections for a total of 21 days. The only detectable drug effects were slight transient ataxia observed occasionally in the high dosage group. No clinical, laboratory, or gross or histopathologic evidence of any butorphanol-related toxicity was encountered in the horses.

## INDICATIONS

Butorphic Injection is indicated for the relief of pain associated with colic in adult horses and yearlings. Clinical studies in the horse have shown that Butorphic Injection alleviates abdominal pain associated with torsion, impaction, intussusception, spasmodic and tympanic colic, and postpartum pain.

## WARNINGS

NOT FOR USE IN HORSES INTENDED FOR FOOD.

NOT FOR HUMAN USE.

## CAUTION

Butorphic Injection, a potent analgesic, should be used with caution with other sedative or analgesic drugs as these are likely to produce additive effects.

There are no well-controlled studies using butorphanol in breeding horses, weanlings, and foals. Therefore, the drug should not be used in these groups.

## ADVERSE REACTIONS

In clinical trials in horses, the most commonly observed side effect was slight ataxia which lasted 3 to 10 minutes. Marked ataxia was reported in 1.5% of the 327 horses treated. Mild sedation was reported in 9% of the horses.

## DOSAGE

The recommended dosage in the horse is 0.1 mg of butorphanol per kilogram of body weight (0.05 mg/lb) by intravenous injection. This is equivalent to 5 mL of Butorphic Injection for each 1000 lbs body weight. The dose may be repeated within 3 to 4 hours but treatment should not exceed 48 hours. Pre-clinical model studies and clinical field trials in horses demonstrate that the analgesic effects of Butorphic Injection are seen within 15 minutes following injection and persist for about 4 hours.

## How Supplied

20 mL multi-dose vials.

## STORAGE

Store at controlled room temperature 15° to 30° C (59° to 86° F).

## REFERENCES

- Pircio AW, Glyls JA, Cavanagh RL, et al. The pharmacology of butorphanol, a 3,14-dihydroxymorphinan narcotic antagonist analgesic. *Arch Int Pharmacodyn Ther* 1976;220:231-257.
- Dobkin AB, Eamkaow S, Caruso FS. Butorphanol and pentazocine in patients with severe postoperative pain. *Clin Pharmacol Ther* 1975;18:547-553.
- Gilbert MS, Hanover RM, Moylan DS, et al. Intramuscular butorphanol and meperidine in postoperative pain. *Clin Pharmacol Ther* 1976;20:359-364.
- Cavanagh RL, Glyls JA, Bierwagen ME. Antitussive properties of butorphanol. *Arch Int Pharmacodyn Ther* 1976;220:258-268.
- Schurig JE, Cavanagh RL, Buyniski JP. Effect of butorphanol and morphine on pulmonary mechanics, arterial blood pressure and venous plasma histamine in the anesthetized dog. *Arch Int Pharmacodyn Ther* 1978;233:296-304.
- Nagashima H, Karamanian A, Malovany R, et al. Respiratory and circulatory effects of intravenous butorphanol and morphine. *Clin Pharmacol Ther* 1976;19:738-745.
- Popio KA, Jackson DH, Ross AM, et al. Hemodynamic and respiratory effects of morphine and butorphanol. *Clin Pharmacol Ther* 1978;23:281-287.
- Robertson JT, Muir WW, Sams R. Cardiopulmonary effects of butorphanol tartrate in horses. *Am J Vet Res* 1981;42:41-44.
- Kalpravidh M, Lumb WV, Wright M, et al. Effects of butorphanol, flunixin, levorphanol, morphine, and xylazine in ponies. *Am J Vet Res* 1984;45:217-223.

Manufactured for

**LLOYD** LLOYD, Inc. of Iowa  
P.O. Box 130  
Shenandoah, IA 51601 USA  
1-800-831-0004  
www.lloydinc.com